

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN8: Among students who rode a bicycle during the past 12 months, the percentage who never or rarely wore a bicycle helmet	88.4 (84.2-91.7)	765	88.4 (74.5-95.2)	61			●
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	8.9 (6.8-11.7)	1,269	17.2 (11.6-25.0)	114		●	
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	7.7 (5.8-10.1)	832	15.2 (7.1-29.5)	73			●
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	15.9 (13.6-18.5)	1,283	22.7 (15.9-31.3)	132			●
QN14: Percentage of students who carried a gun on one or more of the past 30 days	3.9 (2.8-5.4)	1,298	11.2 (6.2-19.5)	133		●	
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	3.7 (2.5-5.6)	1,306	10.4 (4.5-21.9)	133			●
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	7.0 (5.4-9.1)	1,314	14.3 (8.9-22.2)	135		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	7.1 (5.1-9.8)	1,308	21.4 (13.4-32.4)	135		●	
QN18: Percentage of students who were in a physical fight one or more times during the past 12 months	22.3 (19.3-25.6)	1,299	35.4 (26.3-45.7)	132		●	
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	6.9 (5.2-9.1)	1,299	23.1 (14.6-34.6)	132		●	
QN21: Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to	8.0 (6.6-9.6)	1,304	30.1 (24.2-36.7)	134		●	
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	33.7 (30.9-36.7)	1,307	53.6 (42.4-64.5)	133		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	16.0 (14.2-18.0)	1,306	43.7 (31.4-56.8)	134		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	13.7 (11.8-15.7)	1,307	42.5 (32.5-53.1)	130		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN29: Percentage of students who actually attempted suicide one or more times during the past 12 months	7.2 (5.9-8.8)	1,165	35.8 (26.2-46.7)	127		●	
QN30: Percentage of students who made a suicide attempt during the past 12 months that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse	2.7 (2.0-3.5)	1,159	15.0 (9.2-23.4)	122		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN31: Percentage of students who ever tried cigarette smoking, even one or two puffs	41.5 (36.4-46.7)	1,247	62.0 (52.1-70.9)	111		●	
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	6.3 (4.7-8.6)	1,275	21.8 (12.0-36.3)	127		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	12.0 (9.5-15.1)	1,251	35.2 (25.5-46.5)	117		●	
QNFRDIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	2.9 (2.0-4.2)	1,251	18.4 (10.7-29.8)	117		●	
QN34: Among students who reported current cigarette use, the percentage who smoked more than 10 cigarettes per day on the days they smoked during the past 30 days	3.5 (1.3-9.3)	147	24.3 (10.4-46.9)	39		●	
QN35: Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	7.9 (4.6-13.2)	106	-	28			
QN36: Percentage of students who smoked cigarettes on school property on one or more of the past 30 days	2.0 (1.2-3.1)	1,294	11.6 (5.0-24.4)	129		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN38: Among students who reported current cigarette use, the percentage who did not ever try to quit smoking cigarettes during the past 12 months	51.1 (41.5-60.7)	142	49.5 (34.2-64.9)	38			●
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	5.3 (3.7-7.6)	1,311	11.7 (5.9-21.9)	134			●
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	10.4 (7.8-13.7)	1,311	22.8 (15.4-32.6)	133		●	
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	17.4 (13.7-22.0)	1,245	40.7 (31.2-50.9)	114		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	17.8 (15.4-20.5)	1,304	30.3 (23.9-37.6)	135		●	
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	34.8 (29.9-39.9)	1,243	53.9 (42.6-64.8)	119		●	
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	19.7 (16.3-23.5)	1,280	26.7 (18.8-36.3)	126			●
QN46: Among students who reported current alcohol use, the percentage who usually got the alcohol they drank from someone who gave it to them during the past 30 days	37.2 (31.9-42.9)	410	27.2 (15.3-43.7)	59			●
QN47: Percentage of students who used marijuana one or more times during their life	42.2 (36.7-47.9)	1,259	63.5 (48.1-76.6)	119		●	
QN48: Percentage of students who tried marijuana for the first time before age 13 years	8.9 (7.0-11.3)	1,276	16.8 (9.5-27.8)	120			●
QN49: Percentage of students who used marijuana one or more times during the past 30 days	22.2 (18.4-26.5)	1,284	34.1 (24.4-45.4)	126		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	8.0 (6.4-10.1)	1,315	19.7 (11.1-32.4)	135		●	
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	8.4 (6.3-11.0)	1,273	25.4 (16.7-36.8)	114		●	
QN52: Percentage of students who used heroin one or more times during their life	3.2 (2.0-5.1)	1,316	13.3 (7.2-23.1)	134		●	
QN53: Percentage of students who used methamphetamines one or more times during their life	2.9 (1.8-4.7)	1,288	19.3 (11.5-30.5)	131		●	
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	3.7 (2.5-5.4)	1,310	20.1 (10.5-35.2)	134		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	28.7 (25.9-31.8)	1,301	48.8 (37.2-60.5)	127		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	44.3 (38.4-50.4)	1,195	70.3 (60.9-78.3)	103		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	4.0 (2.6-6.0)	1,192	22.1 (12.0-36.9)	104		●	
QN61: Percentage of students who had sexual intercourse with four or more people during their life	9.3 (6.9-12.5)	1,187	29.0 (18.9-41.6)	104		●	
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	30.7 (26.4-35.4)	1,186	61.2 (53.2-68.6)	104		●	
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	20.0 (15.2-25.7)	365	26.1 (15.5-40.6)	62			●
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	41.9 (36.3-47.8)	360	62.0 (49.6-73.0)	62		●	
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	80.8 (76.2-84.7)	359	92.3 (80.3-97.2)	60		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	92.4 (87.9-95.3)	359	97.0 (87.7-99.3)	60		●	
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	73.2 (68.0-77.8)	359	89.2 (77.3-95.3)	60		●	
QNDUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	91.9 (88.2-94.5)	355	97.0 (87.6-99.3)	60		●	
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	13.0 (8.7-18.9)	359	45.9 (27.3-65.8)	60		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

# 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

## Arizona High School Survey Risk Behaviors and Sexual Identity Report

### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) <sup>§</sup>	12.3 (10.6-14.3)	1,243	11.6 (6.7-19.5)	128			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) <sup>§</sup>	9.5 (7.6-11.8)	1,243	12.7 (7.0-22.0)	128			●
QN66: Percentage of students who described themselves as slightly or very overweight	25.3 (23.3-27.3)	1,308	29.3 (20.5-39.9)	133			●
QN68: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight during the past 30 days	14.0 (11.7-16.6)	1,304	31.8 (18.8-48.5)	129		●	
QN69: Percentage of students who took any diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight during the past 30 days	6.5 (5.2-8.1)	1,301	15.7 (7.4-30.3)	127			●
QN70: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight during the past 30 days	6.9 (5.3-9.0)	1,293	25.8 (14.9-40.9)	124		●	
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	18.2 (14.8-22.2)	1,262	31.3 (19.0-47.0)	114			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

<sup>§</sup>Based on reference data from the 2000 CDC Growth Charts.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	71.9 (69.8-73.9)	1,262	77.0 (68.6-83.6)	114			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	11.4 (9.1-14.1)	1,262	21.2 (10.6-37.8)	114			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	4.7 (3.0-7.1)	1,262	14.1 (7.0-26.6)	114		●	
QN78: Percentage of students who did not drink milk or who drank less than three glasses per day of milk during the past seven days	90.4 (88.5-92.0)	1,257	92.5 (87.1-95.8)	112			●
QNMILK0: Percentage of students who drank 0 glasses per day of milk during the past seven days	22.0 (19.3-24.9)	1,257	29.0 (20.8-38.8)	112			●
QNMILK1: Percentage of students who drank less than one glass per day of milk during the past seven days	65.7 (62.9-68.5)	1,257	71.6 (62.1-79.6)	112			●
QNMILK2: Percentage of students who drank less than two glasses per day of milk during the past seven days	80.1 (77.6-82.5)	1,257	83.6 (78.3-87.8)	112			●
QN79: Percentage of students who did not eat breakfast on all of the past seven days	66.8 (63.4-70.0)	1,259	82.7 (71.8-90.0)	112		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNNOBKFT: Percentage of students who ate breakfast on none of the past seven days	14.7 (12.1-17.7)	1,259	31.2 (19.3-46.3)	112		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	56.1 (53.7-58.5)	1,294	75.2 (64.1-83.8)	133		●	
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	16.5 (14.0-19.2)	1,294	21.5 (15.7-28.7)	133			●
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	77.2 (74.3-79.8)	1,294	85.9 (76.6-91.9)	133		●	
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	26.4 (23.4-29.6)	1,288	33.7 (22.2-47.6)	133			●
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	36.2 (31.8-40.9)	1,295	41.2 (31.3-51.8)	132			●
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	60.0 (52.8-66.7)	1,301	62.3 (49.0-73.9)	132			●
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	76.3 (67.3-83.4)	1,301	83.2 (73.7-89.7)	132			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN84: Percentage of students who did not play on sports teams during the past 12 months	48.2 (44.4-52.0)	1,293	57.4 (45.5-68.5)	131			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Other

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	22.2 (19.8-24.9)	1,260	34.9 (26.6-44.3)	112		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who would not describe their grades in school as mostly A's or mostly B's during the past 12 months	33.3 (28.3-38.8)	1,305	46.6 (35.3-58.1)	134		●	
QN88: Percentage of students who rode in a car or other vehicle driven by a friend who had been drinking alcohol one or more times during the past 30 days	17.1 (14.2-20.3)	1,313	26.6 (20.7-33.5)	135		●	
QN89: Percentage of students who carried a gun on school property on one or more of the past 30 days	1.1 (0.6-2.2)	1,304	8.1 (3.4-18.3)	134		●	
QN90: Percentage of students who sometimes, rarely, or never feel safe and secure at school	21.4 (18.8-24.3)	1,312	32.5 (24.5-41.7)	134		●	
QN91: Percentage of students who were ever hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend during the past 12 months	8.2 (6.7-10.0)	1,302	19.6 (13.1-28.4)	132		●	
QN92: Percentage of students who have been harassed or bullied once or twice, monthly, weekly, or daily on school property during the past 12 months	26.0 (22.7-29.7)	1,316	45.1 (32.3-58.5)	135		●	
QN93: Percentage of students who have harassed or bullied someone else once or twice, monthly, weekly, or daily on school property during the past 12 months	18.5 (15.3-22.2)	1,306	31.1 (19.1-46.2)	135		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN94: Percentage of students who have been electronically bullied, such as through e-mail, chat rooms, instant messaging, web sites, or text messaging once or twice, monthly, weekly, or daily during the past 12 months	17.2 (14.6-20.2)	1,312	30.1 (24.0-36.9)	135		●	
QN95: Percentage of students who have electronically bullied someone else, such as through e-mail, chat rooms, instant messaging, web sites, or text messaging once or twice, monthly, weekly, or daily during the past 12 months	9.2 (7.1-11.8)	1,313	26.2 (16.5-39.1)	135		●	
QN96: Percentage of students whose property such as their car, clothing, or books had been stolen or deliberately damaged on school property once or twice, monthly, weekly, or daily during the past 12 months	21.7 (18.5-25.2)	1,299	36.0 (23.7-50.3)	135		●	
QN97: Percentage of students who did something to purposely hurt themselves without wanting to die, such as cutting or burning themselves on purpose, one or more times during the past 12 months	18.1 (15.8-20.6)	1,312	50.8 (40.6-61.0)	134		●	
QN98: Percentage of students who have ever used chewing tobacco, snuff, or dip, such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, or Copenhagen	12.0 (9.7-14.9)	1,301	21.4 (11.3-36.9)	126			●
QN99: Percentage of students who used chewing tobacco, snuff, or dip on school property on one or more of the past 30 days	2.8 (1.9-4.0)	1,302	10.2 (4.6-21.5)	129			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN100: Percentage of students who have ever used a waterpipe (also called a hookah, shisha, or narghile) to smoke tobacco, even one or two puffs	38.8 (34.4-43.5)	1,241	55.9 (48.5-63.1)	107		●	
QN101: Percentage of students who used a waterpipe (also called a hookah, shisha, or narghile) to smoke tobacco on one or more of the past 30 days	21.2 (17.7-25.2)	1,288	37.5 (26.6-49.9)	130		●	
QN102: Percentage of students who were in the same room with someone who was smoking cigarettes on one or more of the past seven days	37.3 (34.0-40.9)	1,312	54.3 (41.5-66.6)	134		●	
QN103: Percentage of students who described the rules about smoking cigarettes inside the home where they live as allowed inside their home or allowed only at some times or in some places in their home	10.8 (8.1-14.1)	1,265	24.4 (14.8-37.5)	112		●	
QN104: Percentage of students who had at least one drink of alcohol on school property on one or more of the past 30 days	4.6 (3.3-6.4)	1,311	19.2 (12.0-29.4)	134		●	
QN105: Percentage of students who used marijuana on school property one or more times during the past 30 days	4.5 (3.1-6.6)	1,306	15.1 (8.3-25.9)	133		●	
QN106: Percentage of students who used any form of cocaine, including powder, crack, or freebase, one or more times during the past 30 days	4.0 (2.9-5.5)	1,311	17.2 (8.9-30.6)	135		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN107: Percentage of students who have sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during the past 30 days	2.8 (1.5-5.1)	1,312	14.0 (7.4-25.1)	135		●	
QN108: Percentage of students who have used methamphetamines (also called speed, crystal, crank, or ice) one or more times during the past 30 days	2.3 (1.1-4.8)	1,271	10.0 (5.3-18.2)	115		●	
QN109: Percentage of students who have used ecstasy (also called MDMA) one or more times during the past 30 days	4.6 (2.9-7.5)	1,313	18.7 (11.8-28.2)	135		●	
QN110: Percentage of students who have attended school under the influence of alcohol or other illegal drugs, such as marijuana or cocaine, one or more times during the past 12 months	17.8 (15.2-20.8)	1,310	33.4 (21.8-47.4)	135		●	
QN111: Percentage of students who have taken over-the-counter drugs to get high one or more times during the past 30 days	6.4 (5.2-7.9)	1,311	25.7 (16.3-38.0)	135		●	
QN112: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during the past 30 days	8.6 (6.7-10.8)	1,314	22.7 (14.7-33.4)	135		●	
QN113: Percentage of students whose first sexual intercourse partner was three or more years older	5.9 (4.8-7.3)	1,305	17.1 (11.0-25.4)	135		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Arizona High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN114: Percentage of students who are not sure if their parents or other adults had ever talked with them or whose parents or other adults in their family have not ever talked with them about what they expect them to do or not to do when it comes to sex	30.4 (27.3-33.6)	1,266	36.7 (28.8-45.3)	113			●
QN117: Percentage of students who ate fruit or drank 100% fruit juice less than one time per day during the past seven days	70.8 (68.1-73.4)	1,257	68.7 (60.3-76.0)	115			●
QN118: Percentage of students who ate vegetables, such as green salads, carrots, broccoli, or potatoes, less than one time per day during the past seven days	66.7 (62.9-70.4)	1,264	70.0 (59.0-79.1)	114			●
QN119: Percentage of students who had been told by a doctor or nurse that they had asthma and who still have asthma	8.9 (7.1-11.1)	1,255	19.5 (13.9-26.8)	111		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.