

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	7.5 (5.7-9.7)	1,752	13.4 (7.9-21.9)	245			●
QN9: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle, one or more times during the 30 days before the survey)	18.7 (15.4-22.6)	1,749	24.6 (15.7-36.3)	243			●
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	5.7 (4.3-7.6)	985	9.3 (5.0-16.8)	121			●
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	15.1 (11.8-19.0)	1,731	19.2 (10.8-31.8)	241			●
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	3.0 (2.1-4.4)	1,740	5.7 (2.8-11.3)	241			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)	6.1 (4.4-8.3)	1,751	7.7 (3.9-14.7)	241			●
QN15: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	8.1 (5.7-11.4)	1,749	20.2 (14.1-28.0)	244		●	
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)	7.4 (5.3-10.3)	1,762	10.1 (6.5-15.4)	244			●
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	19.4 (16.4-22.6)	1,743	31.4 (26.4-36.9)	240		●	
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	5.0 (3.8-6.6)	1,717	10.6 (6.5-16.8)	239		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN19: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	6.1 (4.8-7.7)	1,750	23.4 (14.9-34.6)	243		●	
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)	15.4 (13.4-17.8)	1,734	40.9 (31.6-51.0)	237		●	
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)	12.7 (10.6-15.1)	1,754	32.3 (22.7-43.8)	242		●	
QN25: Percentage of students who felt sad or hopeless (almost every day for ≥2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	31.6 (28.7-34.7)	1,747	69.7 (58.9-78.8)	239		●	
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)	15.1 (13.0-17.6)	1,746	49.9 (38.9-60.8)	240		●	
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	11.6 (10.0-13.3)	1,753	37.8 (30.3-46.0)	238		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey

Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	8.2 (6.5-10.3)	1,483	34.7 (25.3-45.4)	211		●	
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	2.5 (1.6-3.7)	1,464	20.5 (12.6-31.4)	210		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)	26.9 (22.3-32.0)	1,699	50.4 (37.2-63.6)	232		●	
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)	7.2 (5.4-9.4)	1,703	15.8 (9.2-25.6)	231		●	
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	5.4 (4.0-7.2)	1,690	19.4 (12.6-28.7)	224		●	
QNFRDIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)	1.3 (0.8-2.1)	1,690	4.0 (1.2-12.5)	224			●
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	1.0 (0.6-1.7)	1,690	1.1 (0.3-4.8)	224			●
QN33: Percentage of students who smoked more than 10 cigarettes per day (on the days they smoked during the 30 days before the survey, among students who currently smoked cigarettes)	7.9 (2.8-20.4)	88	2.9 (0.4-20.3)	35			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])	49.6 (44.2-55.0)	1,737	64.4 (55.0-72.9)	241		●	
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)	14.3 (10.5-19.1)	1,490	30.8 (21.9-41.4)	210		●	
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, Copenhagen, Camel Snus, Marlboro Snus, General Snus, Ariva, Stonewall, or Camel Orbs], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)	4.0 (2.4-6.4)	1,700	7.3 (3.8-13.8)	237			●
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)	4.7 (3.1-7.0)	1,730	10.9 (6.3-18.2)	238		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	7.7 (5.7-10.5)	1,673	21.5 (14.4-30.8)	226		●	
QNTB3: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco (on at least 1 day during the 30 days before the survey)	10.4 (8.2-13.1)	1,637	24.9 (16.6-35.6)	225		●	
QNTB4: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco or electronic vapor products (on at least 1 day during the 30 days before the survey)	18.9 (14.9-23.6)	1,415	41.5 (31.5-52.2)	202		●	
QN39: Percentage of students who did not try to quit using all tobacco products (including cigarettes, cigars, smokeless tobacco, shisha or hookah tobacco, and electronic vapor products, ever during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)	55.8 (42.8-68.0)	299	50.9 (39.2-62.5)	75			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)	16.6 (13.8-19.8)	1,711	27.5 (19.0-38.1)	233		●	
QN42: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	30.8 (26.9-35.1)	1,617	52.7 (40.7-64.3)	213		●	
QN43: Percentage of students who usually got the alcohol they drank by someone giving it to them (during the 30 days before the survey, among students who currently drank alcohol)	41.2 (36.1-46.5)	466	28.3 (18.8-40.3)	95	●		
QN44: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)	16.5 (13.5-20.1)	1,713	31.9 (22.3-43.2)	234		●	
QN47: Percentage of students who tried marijuana for the first time before age 13 years	6.2 (4.8-8.1)	1,749	14.4 (8.7-22.7)	236		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN48: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	17.2 (13.9-21.3)	1,748	37.7 (26.7-50.1)	238		●	
QN49: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	5.0 (3.7-6.9)	1,760	9.2 (4.6-17.4)	243			●
QN51: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	0.9 (0.5-1.7)	1,760	6.9 (2.6-16.8)	245			●
QN52: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	1.4 (0.9-2.1)	1,760	7.2 (3.0-16.4)	242			●
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)	3.1 (2.1-4.5)	1,762	10.2 (5.4-18.3)	243		●	
QN56: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their	13.3 (11.4-15.5)	1,757	30.7 (22.8-40.0)	239		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN58: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	27.3 (24.3-30.5)	1,750	45.4 (33.3-58.0)	238		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	32.7 (28.9-36.8)	1,589	46.3 (35.7-57.3)	196		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	2.4 (1.3-4.7)	1,588	5.4 (2.3-12.2)	198			●
QN61: Percentage of students who had sexual intercourse with four or more persons during their life	7.9 (6.2-10.2)	1,587	11.0 (6.3-18.7)	197			●
QN62: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	24.1 (20.6-28.1)	1,585	36.8 (27.7-46.9)	197		●	
QN63: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	15.2 (10.4-21.7)	356	34.4 (15.1-60.6)	63			●
QN64: Percentage of students who did not use a condom during last sexual intercourse (among students who were currently sexually active)	47.7 (38.1-57.5)	352	46.4 (28.2-65.5)	62			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN65: Percentage of students who did not use birth control pills before last sexual intercourse (to prevent pregnancy among students who were currently sexually active)	79.5 (68.5-87.4)	344	85.1 (68.6-93.7)	62			●
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	96.1 (91.2-98.3)	344	98.3 (87.3-99.8)	62			●
QNSHPARG: Percentage of students who did not use a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	97.1 (93.1-98.8)	344	96.4 (85.7-99.2)	62			●
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	72.7 (61.7-81.5)	344	79.7 (62.7-90.2)	62			●
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	88.8 (80.8-93.7)	343	90.2 (74.8-96.6)	62			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey

Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	14.3 (9.1-21.8)	344	30.3 (16.6-48.7)	62			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]	11.4 (9.3-13.7)	1,629	17.5 (11.2-26.5)	219			●
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]	15.2 (12.5-18.4)	1,629	18.3 (13.4-24.5)	219			●
QN68: Percentage of students who described themselves as slightly or very overweight	29.5 (24.7-34.7)	1,744	44.9 (39.0-51.0)	240		●	
QN69: Percentage of students who were not trying to lose weight	56.0 (50.5-61.4)	1,744	31.0 (24.7-38.1)	243	●		
QN70: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	32.3 (28.9-35.8)	1,744	33.8 (23.8-45.5)	242			●
QN71: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	13.8 (11.7-16.2)	1,732	18.9 (12.8-27.1)	240			●

*Confidence interval.

[†]Based on t-test analyses, $p < 0.05$.

[§]Based on reference data from the 2000 CDC Growth Charts. In 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	8.0 (6.5-10.0)	1,726	10.7 (5.6-19.5)	240			●
QN72: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)	39.6 (35.5-43.8)	1,739	44.4 (34.7-54.5)	243			●
QN73: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)	35.2 (32.2-38.2)	1,738	39.2 (31.4-47.7)	242			●
QN74: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)	51.4 (48.4-54.4)	1,727	56.7 (47.6-65.5)	241			●
QN75: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)	18.3 (15.6-21.4)	1,734	22.8 (16.9-30.1)	242			●
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	6.4 (5.0-8.2)	1,712	11.5 (6.8-18.8)	239			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	16.9 (14.0-20.4)	1,728	19.6 (12.3-29.7)	243			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	9.2 (7.1-11.9)	1,728	14.5 (7.8-25.2)	243			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	3.8 (2.5-5.6)	1,728	6.0 (3.3-10.4)	243			●
QN77: Percentage of students who did not drink milk (during the 7 days before the survey)	24.8 (22.0-28.0)	1,710	30.7 (21.0-42.5)	238			●
QN78: Percentage of students who did not eat breakfast (during the 7 days before the survey)	15.9 (14.4-17.4)	1,721	28.2 (20.0-38.2)	243		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey

Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNBK7DAY: Percentage of students who did not eat breakfast on all 7 days (during the 7 days before the survey)	63.3 (59.5-66.9)	1,721	79.0 (71.4-85.0)	243		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN79: Percentage of students who were not physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	51.2 (47.2-55.2)	1,730	67.3 (59.1-74.5)	236		●	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	15.6 (12.7-19.1)	1,730	21.9 (17.0-27.9)	236		●	
QNPA7DAY: Percentage of students who were not physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	73.8 (70.5-76.8)	1,730	84.1 (77.2-89.3)	236		●	
QN80: Percentage of students who watched television 3 or more hours per day (on an average school day)	19.0 (15.9-22.4)	1,711	19.9 (12.5-30.2)	238			●
QN81: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)	37.3 (32.6-42.3)	1,731	48.1 (39.1-57.3)	238		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN82: Percentage of students who did not attend physical education (PE) classes on 1 or more days (in an average week when they were in school)	52.5 (43.1-61.7)	1,717	61.7 (47.9-73.8)	236			●
QNDLYPE: Percentage of students who did not attend physical education classes on all 5 days (in an average week when they were in school)	62.1 (54.0-69.5)	1,717	73.2 (61.1-82.6)	236		●	
QN83: Percentage of students who did not play on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)	46.3 (41.2-51.4)	1,704	59.7 (48.9-69.7)	235		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey

Risk Behaviors and Sexual Identity Report

Other

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN89: Percentage of students who described their grades in school as mostly C's, D's, or F's (during the 12 months before the survey)	30.0 (23.9-36.8)	1,700	32.3 (25.2-40.3)	232			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN90: Percentage of students who carried a gun on school property (on at least 1 day during the 30 days before the survey)	1.7 (0.7-3.7)	1,592	2.9 (1.1-7.4)	222			●
QN91: Percentage of students who sometimes, rarely, or never feel safe and secure at school	20.9 (15.2-28.0)	1,580	42.2 (31.4-53.9)	221		●	
QN92: Percentage of students who were ever hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend (during the 12 months before the survey)	8.2 (6.4-10.5)	1,584	19.7 (13.4-28.0)	220		●	
QN93: Percentage of students who bullied someone on school property (during the 12 months before the survey)	7.4 (5.8-9.3)	1,588	21.4 (13.8-31.5)	217		●	
QN94: Percentage of students who electronically bullied someone (counting bullying through texting, Instagram, Facebook, or other social media, during the 12 months before the survey)	5.8 (4.3-7.8)	1,580	14.5 (8.8-23.1)	218		●	
QN95: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)	16.6 (13.7-20.1)	1,588	54.5 (46.1-62.6)	215		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who were in the same room with someone who was smoking cigarettes (on at least 1 day during the 7 days before the survey)	30.3 (26.0-34.9)	1,582	47.6 (38.1-57.4)	220		●	
QN97: Percentage of students who described the rule about smoking cigarettes inside the home where they live as always allowed inside their home or allowed only at some times or in some places inside their home	6.0 (4.5-8.1)	1,579	16.6 (10.8-24.7)	217		●	
QN98: Percentage of students who currently used a waterpipe to smoke tobacco (also called a hookah, shisha, or narghile on at least 1 day during the 30 days before the survey)	5.8 (4.5-7.6)	1,571	15.8 (8.4-27.7)	217		●	
QN99: Percentage of students who ever used chewing tobacco, snuff, dip, snus, or dissolvable tobacco products (such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, Copenhagen, Camel Snus, Marlboro Snus, General Snus, Ariva, Stonewall, or Camel Orbs, not counting any electronic vapor products)	10.9 (8.3-14.3)	1,586	12.7 (6.9-22.0)	217			●
QN100: Percentage of students who currently had at least one drink of alcohol on school property (on at least 1 day during the 30 days before the survey)	4.0 (2.9-5.4)	1,589	13.0 (7.1-22.6)	221		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN101: Percentage of students who used marijuana on school property (one or more times during the 30 days before the survey)	4.8 (3.1-7.3)	1,577	12.8 (7.6-20.8)	221		●	
QN102: Percentage of students who currently used any form of cocaine (including powder, crack, or freebase, one or more times during the 30 days before the survey)	2.4 (1.8-3.3)	1,586	9.1 (4.9-16.3)	220		●	
QN103: Percentage of students who currently sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high (one or more times during the 30 days before the survey)	2.4 (1.6-3.7)	1,550	11.9 (6.3-21.5)	212		●	
QN104: Percentage of students who currently used methamphetamines (also called “speed,” “crystal,” “crank,” or “ice,” one or more times during the 30 days before the survey)	1.0 (0.5-1.9)	1,583	4.5 (2.0-10.2)	219			●
QN105: Percentage of students who currently used ecstasy (also called “MDMA,” one or more times during the 30 days before the survey)	2.0 (1.3-3.2)	1,588	6.6 (2.9-14.0)	222			●
QN106: Percentage of students who currently took over-the-counter drugs to get high (one or more times during the 30 days before the survey)	4.0 (3.2-5.0)	1,588	18.0 (12.2-25.8)	221		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN107: Percentage of students who attended school under the influence of alcohol or other illegal drugs (such as marijuana or cocaine, one or more times during the 12 months before the survey)	12.8 (10.7-15.1)	1,579	35.4 (27.1-44.7)	219		●	
QN108: Percentage of students who reported their partners were three or more years older than themselves the first time they had sexual intercourse (among students who have had sexual intercourse)	11.2 (8.2-15.2)	503	23.6 (16.2-33.0)	104		●	
QN109: Percentage of students who reported their parents or other adults in their family have not ever talked with them about what they expect them to do or not to do when it comes to sex or who were not sure if their parents or other adults had ever talked with them	33.4 (29.9-37.1)	1,576	36.7 (25.2-50.0)	214			●
QN111: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight (also called fasting, during the 30 days before the survey)	10.8 (8.6-13.4)	1,572	41.2 (33.0-49.9)	211		●	
QN112: Percentage of students who took some diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight (during the 30 days before the survey)	5.5 (4.0-7.5)	1,565	18.5 (11.5-28.4)	211		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2017 YOUTH RISK BEHAVIOR SURVEY RESULTS

Arizona (Including Charter Schools) High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN113: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	5.3 (4.1-6.8)	1,563	26.5 (18.3-36.7)	213		●	
QN114: Percentage of students who did not see a doctor or nurse (for a check-up or physical exam when they were not sick or injured during the 12 months before the survey)	36.6 (32.3-41.1)	1,558	37.7 (30.9-45.0)	216			●
QN115: Percentage of students who have physical disabilities or long-term health problems (meaning 6 months or more)	13.9 (11.9-16.2)	1,551	34.2 (28.3-40.6)	214		●	
QN116: Percentage of students who have long-term emotional problems or learning disabilities (meaning 6 months or more)	13.7 (11.4-16.4)	1,543	44.9 (33.7-56.6)	211		●	
QN117: Percentage of students who did not go to school because they were sick or in pain (at least one day during the 30 days before the survey)	46.2 (43.3-49.1)	1,557	60.6 (52.2-68.4)	213		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.